



OPEN

European primary forest database v2.0

DATA DESCRIPTOR

Francesco Maria Sabatini *et al.*[#]

Primary forests, defined here as forests where the signs of human impacts, if any, are strongly blurred due to decades without forest management, are scarce in Europe and continue to disappear. Despite these losses, we know little about where these forests occur. Here, we present a comprehensive geodatabase and map of Europe's known primary forests. Our geodatabase harmonizes 48 different, mostly field-based datasets of primary forests, and contains 18,411 individual patches (41.1 Mha) spread across 33 countries. When available, we provide information on each patch (name, location, naturalness, extent and dominant tree species) and the surrounding landscape (biogeographical regions, protection status, potential natural vegetation, current forest extent). Using Landsat satellite-image time series (1985–2018) we checked each patch for possible disturbance events since primary forests were identified, resulting in 94% of patches free of significant disturbances in the last 30 years. Although knowledge gaps remain, ours is the most comprehensive dataset on primary forests in Europe, and will be useful for ecological studies, and conservation planning to safeguard these unique forests.

Background & Summary

The importance of primary forests is widely recognized^{1,2}, they provide refuge to forest biodiversity³, and act as a buffer to species loss in human-dominated landscapes⁴. Second, primary forests play an important role in climate change mitigation. At the local scale, they buffer the adverse effects of increasing temperature on understory biodiversity, as they often have cooler forest-floor summer temperatures compared to secondary forests⁵. At the global scale they contribute to climate stability by storing large quantities of carbon, both in the biomass and in soils^{1,6,7}. Third, primary forests often serve as a reference for developing close-to-nature forest management, or for benchmarking restoration efforts⁸. Finally, these forests are an irreplaceable part of our natural heritage, shape the cultural identities of local communities, and have a high intrinsic value⁹.

In Europe, as in many human-dominated regions, most forests are currently managed¹⁰, often with increasing harvest intensities^{11,12}. As a result, despite the general trend of increasing total forest area, primary forests are scarce and continue to disappear¹³. For instance, Romania hosts some of the largest swaths of primary forest in Central Europe and faced a sharp increase in logging rates since 2000. This has resulted in significant primary forest loss, even within protected areas^{13–15}. In Poland, the iconic Białowieża Forest was recently in the spotlight after the controversial decision from the Polish National Forest Holding, now nullified by the Court of Justice of the European Union¹⁶, to implement salvage logging followed by tree planting after a bark beetle outbreak¹⁷. Widespread loss of primary forests also occurred in Ukraine¹⁸, Slovakia¹⁹, or in the boreal North, e.g., in the Russian North-West, where 4.6 Mha of primary forest were lost since 2001^{13,20}. Effective protection of Europe's primary forests is therefore urgently needed²¹.

In the newly released 'Biodiversity Strategy for 2030', the European Commission emphasized the need to define, map, monitor and strictly protect all of the EU's remaining primary forests². Reaching these objectives requires complete and up-to-date data on primary forests' location and protection status. Such data could inform both policy making and conservation planning, as well as research, for instance by highlighting areas where primary forests are either scarce, or poorly studied. Yet, many data gaps remain on the location and conservation status of EU's primary forests^{21,22}. Only a few countries conducted systematic, on-the-ground inventories^{19,23}. For most countries data are either only available for a few well-studied forests^{24–26}, or are limited to the distribution of potential (=unconfirmed) primary forests, typically predicted statistically or via remote sensing^{27–29}. Despite past

[#]A full list of authors and their affiliations appears at the end of the paper.




Fig. 1 Overview of the primary forest patches contained in the EPFD v2.0. Both points and polygons were magnified to improve visibility.

efforts for harmonizing data^{24,30}, only recently has the first map of primary forests been released for Europe^{31,32} together with a first assessment of their conservation status²¹.

In a previous effort, we assembled a first European Primary Forest database (EPFD v1.0) that included 32 local-to-national datasets, plus data from a literature review and a survey, resulting in the mapping of a total of ~1.4 Mha of primary forest³¹. This was only about one fifth of the estimated 7.3 Mha of undisturbed forest still occurring in Europe, excluding Russia¹⁰. Also, most of the data collected in our v1.0 database were not open-access, and could thus not be used without the explicit consent of their respective copyright holders.

Here, we build on those efforts to progress further towards a complete map of Europe's primary forests. First, we secured permission from all data holders to release all data with open-access. Second, we aggregated and harmonized 16 additional regional-to-continental spatial datasets to now cover a total of 48 independent datasets. The EPFD v2.0 contains 18,411 non-overlapping primary forest patches (plus 299 point features) covering an area of 41.1 Mha (37.4 Mha in European Russia alone; Fig. 1) across 33 countries (Table 1)³³. Key improvements of this new database include (a) filling major regional gaps, including European Russia, the Balkan Peninsula, the Pyrenees and the Baltic region, (2) mapping potential primary forests for Sweden and Norway (additional 16,311 polygons and 2.5 Mha - Fig. 2), two key regions where complete inventories are currently unavailable, and (3) an update of our literature review to January 2019.

Methods

Primary forest definition. Defining primary forests is controversial, and a range of different definitions have been put forward over the years²². In this paper, as in our previous work, we follow the FAO definition that defines primary forests³⁴ as “naturally regenerated forest of native tree species, where there are no clearly visible indications of human activities and the ecological processes are not significantly disturbed”.

We operationalized this definition using the framework proposed by Buchwald³⁵, where ‘primary forest’ is used as an umbrella term to include forests with different levels of naturalness, such as primeval, virgin, near-virgin, old-growth and long-untouched forests. Based on this framework, a forest qualifies as primary if the signs of former human impacts, if any, are strongly blurred due to decades (at least 60–80 years) after the end of forest management³⁵. This time limit, however, depends on how modified the forest was at the starting point, and only applies in the case of traditional management, such as patch felling, partial coppicing, or selective logging. Stands regenerating naturally after a clear cut would therefore require a longer time period to be considered a primary forest (i.e., 60–80 years plus the length of a typical rotation cycle). Our definition of primary forests, therefore, does not imply that these forests were never cleared or disturbed by humans. We consider this is in line with the Convention of Biological Diversity (CBD, <https://www.cbd.int/forest/definitions.shtml>), acknowledging that the concept of primary forests has a different connotation in Europe than in the rest of the world.

Finally, our collection of primary forests includes mainly old-growth, late-successional forests, but also some early seral stages and young forests that originated after natural disturbances and natural regeneration, without subsequent management. In case of large primary forest tracts (>250 ha), our polygons can also locally include land not covered by trees.

Data collection. To create the EPFD v2.0, we first expanded and updated the literature review on primary forests we had originally carried out for EPFD v1.0³¹, which only considered the period 2000–2017, and excluded

Country	Num. features (Polygons\ Points)	Tot. estimated area (1,000 ha)	Sources (Dataset IDs)
Albania	13\6	13.36	0, 1, 47, 54
Austria	34\2	1.46	9, 35, 49
Belarus	3\0	188.29	46
Bosnia and Herzegovina	4\12	4.1	0, 2, 50, 53
Bulgaria	483\2	56.77	0, 3, 4, 35
Croatia	45\3	6.24	0, 5, 9
Czechia	86\10	9.07*	0, 6, 9
Denmark	0\24	1.68	7
Estonia	0\29	0.05*	0, 8
Finland	1,008\3	2,817.36*	0, 12, 38, 39
France	106\7	10.86*	0, 13, 14, 35, 37
Germany	25\21	13.65*	0, 9, 15, 35
Greece	5\2	1.75*	0, 16
Italy	86\12	6.84*	0, 18, 35, 55
Latvia	3\0	4.79	40
Lithuania	20\0	32.05	19
Moldova	0\1	0.03	35
Montenegro	2\0	2.85	2, 50
Netherlands	3\0	0.08	36
North Macedonia	5\1	0.81	1, 20
Norway	240\1	280.05*	0, 21, 36, 43
Poland	66\5	21.15*	0, 22, 35
Portugal	32\21	15.75*	23, 24
Romania	3,571\6	59.11*	0, 1, 25, 32, 33, 35
Russian Federation	3,082\3	37,417.69*	0, 51
Serbia	14\4	7.78	0, 35, 36, 44, 45
Slovakia	290\4	10.98	0, 9, 26
Slovenia	170\1	9.53	0, 27
Spain	44\58	9.4*	0, 41, 52
Sweden	0\51	32.81*	0, 29, 35
Switzerland	5\5	2.29	0, 30, 35
Ukraine	8,966\3	97.8*	0, 1, 32
United Kingdom	0\2	0.1	9
Total	18,411\299	41,136.53*	

Table 1. Summary of primary forest data across European countries. Dataset IDs correspond to those in Online-only Table 1. * Some point features have no information on forest patch area.

European Russia. Specifically, we added all scientific studies published between January 2000 and January 2019 for Russia, and those published in 2017–2019 for the rest of Europe. We identified relevant publications in the ISI Web of Knowledge using the search terms “(primary OR virgin OR old-growth OR primeval OR intact) AND forest*” in the title field. Based on our own interpretation of commonly used forest terms, we deliberately excluded terms such as “unmanaged” (meaning: not under active management), “ancient” (never cleared for agriculture) or “natural” (stocked with naturally regenerated native trees). These terms indicate conditions that are necessary, but not sufficient for considering a forest as primary. Finally, we refined our search using geographical and subject filters. The literature search returned 129 candidate papers. After screening their content, we added 23 additional primary forest stands (10 in European Russia, 13 in the rest of Europe), from 13 studies (four from European Russia, and nine from the rest of Europe).

Building the EPFD v1.0³¹ involved reaching out to 134 forest experts. For v2.0 we contacted an additional 75 experts with knowledge on forests or forestry, and invited them to add spatially-explicit data on primary forests to our database. We focussed on experts from geographical regions poorly covered in v1.0. We received 56 answers, which led to the incorporation of 16 new datasets in our map. Given the context-dependency of definitions used in regional mapping projects, new datasets were only included if we could find an explicit equivalence between country-specific forest definitions and our definition framework³⁵. This was done after discussing with data contributors the criteria and categories used for constructing their datasets, which we then mapped onto our definition framework. Depending on the datasets, these criteria included: (1) forest age or structural variables^{19,23,36}, (2) legal designation²⁵ or year since onset of protection³⁷, (3) time since last anthropogenic disturbance³⁸, or (4) the lack of human impacts and infrastructures³⁹.

We integrated all data into a geodatabase, which contains primary forests either as polygons (if information on the forest boundary was available) or point locations (when having only an approximate centre location). We set 0.5 hectares as minimum mapping unit, although only a few of the datasets already contained in v1.0 contained




Fig. 2 Overview of the maps of potential primary forests of Sweden and Norway.

polygons smaller than 2 ha (i.e., the minimum mapping unit originally used). If available, we included a set of basic descriptors for each patch: name, location, naturalness level (based on³⁵), extent, dominant tree species, disturbance history and protection status. In total, our map harmonizes 48 regional-to-continental datasets of primary forests (Online-only Table 1). All data is open-access³³, except for three datasets that we kept confidential, either for conservation or copyright reasons. These datasets are: ‘Hungarian Forest Reserve monitoring’ (ID 17, custodian: Ferenc Horváth); ‘Ancient and Primeval Beech Forests of the Carpathians and Other Regions of Europe’^{40,41} (ID 34, copyright: UNESCO), and ‘Potential OGF and primary forest in Austria’ (ID 48, custodian: Matthias Schickhofer). Additional non-open access polygons also exist for the dataset ‘Strict Forest Reserves in Switzerland’ (ID 30, custodian: Jonas Stillhard). These data are here referred to for transparency, but are neither included in the statistics and summaries reported here, nor in any of the remote-sensing analysis below.

Post-processing. To provide common descriptions for all features contained in the geodatabase, we integrated the basic descriptors detailed above with a range of attributes derived by intersecting all polygons or points of primary forests with layers of: 1) biogeographical regions, 2) protected areas, 3) forest type, and 4) forest cover.

We used the map of biogeographical regions⁴² to assign each primary forest point or polygon to one of the following ten classes: 1. Alpine, 2. Arctic, 3. Atlantic, 4. Black Sea, 5. Boreal, 6. Continental, 7. Macaronesia, 8. Mediterranean, 9. Pannonian, 10. Steppic. Similarly, we derived information on protection status and time since onset of protection for each primary forest polygon or point based on the World Database on Protected Areas (WDPA - <https://www.protectedplanet.net>). We simplified the original IUCN classification to three classes: 1. strictly protected – (IUCN category I); 2. protected – (IUCN categories II–VI + not classified); 3. not protected. This is a conservative aggregation recognizing the fact that, in certain contexts, logging and salvage logging are allowed inside national parks, at least in the buffer zone. In case of polygons, we considered a primary forest patch as protected if > 75% of its surface was within a WDPA polygon. When better information on the protection status of a forest patch was available directly from data contributors, we gave priority to this source. We also assigned each primary forest polygon or point to one of the forest categories defined by the European Environmental Agency⁴³. The spatial information was derived by simplifying the map of Potential Vegetation types for Europe⁴⁴, after creating an expert-based cross-link table²¹, which ties together forest categories and potential vegetation types reported in Table 4.1 from⁴³. After excluding forest plantations, the remaining 13 categories comprised: 1. Boreal forest; 2. Hemiboreal forest and nemoral coniferous and mixed broadleaved-coniferous forest; 3. Alpine coniferous forest; 4. Acidophilous oakwood and oak-birch forest; 5. Mesophytic deciduous forest; 6. Lowland to submountainous beech forest; 7. Mountainous beech forest; 8. Thermophilous deciduous forest; 9. Broadleaved




Fig. 3 Workflow and data sources for the map of potential primary forests in Sweden. Data on woodland key habitats derive from⁶⁰ (see also: <https://www.skogsstyrelsen.se>); forest with conservation value from^{61,62}, forest core areas from⁶³, continuity forests from^{64,65}, protected mountain coniferous forests from⁶⁶, clear cuts and fellings from <https://www.skogsstyrelsen.se>.

evergreen forest; 10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions; 11. Mire and swamp forest; 12. Floodplain forest; 13. Non-riverine alder, birch or aspen forest. For each primary forest polygon (but not for points), we reported the two most common forest categories. Finally, we extracted for each primary forest polygon the actual share covered by forest. We did this, because larger primary forest polygons in high naturalness classes can encompass land temporarily or permanently not covered by trees. We used a tree cover density map for the year 2010 for these regions from⁴⁵. All post-processing was performed in R (v3.6.1)⁴⁶.

Potential primary forests of Sweden and Norway. For Sweden and Norway, where abundant geographic information was available on forest distribution, we created maps of potential (but so far unconfirmed) primary forests. For Sweden, we derived a workflow to create a map of potential primary forests as detailed in Fig. 3. This yielded 14,300 polygons covering a total area of 2.4 Mha.

For Norway, even though we were able to include two datasets of confirmed primary forests, additional primary forest is expected to exist. Therefore, we derived a map of potential primary forests, based on the “*Viktige Naturtyper*” dataset from the Norwegian Environment Agency⁴⁷, which maps different habitat types of high conservation value both inside and outside forested areas. We extracted all polygons larger than 10 ha classified as “*old forest types*” (=“*gammelskog*”), i.e., forests that have never been clearcut and are in age classes of 120 years or older. This yielded 2,103 polygons covering a total area of 0.1 Mha.

Importantly, these layers were neither directly integrated in our composite map, nor used to calculate country level statistics as they only represent a first approximation of the primary forest situation in these countries, so far without ground validation. Yet, we included these layers in our geodatabase with the goal of directing future ground-based mapping efforts.

Data Records

The EPFD v2.0³³ is composed of 48 individual datasets (Online-only Table 1) and the two layers of potential primary forests for Sweden and Norway. We integrated the 48 datasets into two composite feature classes, after excluding all duplicated/overlapping polygons across individual datasets.

1) *EU_PrimaryForests_Polygons_OA_v20*

- Composite feature class combining the forest patches classified as “primary forest” based on polygon data sources described in Online-only Table 1
- Data type: Polygon Feature Class

2) *EU_PrimaryForests_Points_OA_v20*

- Composite feature class combining forest locations classified as “primary forest”, based on point data sources described in Online-only Table 1. Only points not overlapping with polygons in (1) reported.
- Data type: Point Feature Class




Fig. 4 Estimation of data completeness. Ratio between the total primary forest area in the EPFD v2.0 and the country estimate of ‘forest undisturbed by man’ (indicator 4.3) from Forest Europe¹⁰ or, if unavailable, the country estimates of primary forests based on FAO’s Forest Resources Assessment⁴⁸. Gray polygons represent countries where Forest Europe (or FAO) reports no forest undisturbed by man (‘No Reported PF’).

The individual datasets are also included in the geodatabase, inside the feature dataset ‘*European_PrimaryForests*’. The whole database is stored in Figshare (<https://doi.org/10.6084/m9.figshare.13194095.v1>)³³. The file format is ESRI personal geodatabase (.mdb). Each feature class in the geodatabase follows the structure described in Online-only Table 2. A full description of each individual dataset is reported in the metadata file ‘*DATASET_overview_v2.0_20201030.docx*’, available at the same link.

Technical Validation

We benchmarked our data against country-level statistics on primary forest extent. Although we had no direct control of the raw data contained in our database, the fact that all our information on primary forest locations derives either from peer-reviewed scientific literature, or was field-checked by trained researchers and/or professionals suggests high data reliability. We made sure to have a common understanding with data contributors about forest definitions [i.e.^{34,35}], and only included a dataset in the EPFD if we could find an explicit equivalence with the forest definitions we used. Additional information on the harmonization process is reported for individual datasets in the metadata accompanying our geodatabase.

An additional, wall-to-wall validation of our database using remotely-sensed information is currently impossible. Remote sensing data only cover the last 35 years, and even if high resolution laser ranging (LIDAR) might become available in the future, at the moment no reliable workflow exists for mapping primary forests from such multi-sensor data. The alternative is field work, which is clearly unfeasible given the huge area covered by our database, the large number of polygons, and the cost and time effort that would be required for a statistically valid ground sample of data. Still, remote sensing data can be helpful for checking whether a patch of primary forest underwent human disturbance after it was delineated and that is why we implemented a semi-automatic procedure based on Landsat satellite-image time series (1985–2018) (see below).

Benchmarking against country-level statistics. Our database contains most of the geographical information currently available on primary forests in Europe, but we do not claim this data is complete. To benchmark the completeness of our map, we calculated the ratio between the area of primary forest in our database at country level, and the estimated area of “forest undisturbed by man” from the indicator 4.3 in the Forest Europe report¹⁰ or, for those countries where this information is not available, from FAO’s Forest Resources Assessment⁴⁸. Although the definition of “forest undisturbed by man” in Forest Europe is consistent with our definition of primary forest¹⁰, it must be noted that these country-level estimates stem from national inventories or other studies, and data quality varies from country to country⁴⁹. The comparison presented here should, therefore, be taken with caution (Fig. 4).

Forest Europe reports no primary forest for some western European countries (Spain, France, Belgium, Netherlands, Germany, United Kingdom and Ireland), although for most of these countries we did find information on at least a handful of primary forest sites. The coverage of our map was also higher than expected for some Eastern European countries (e.g., Ukraine, Belarus, Lithuania), as well as Norway and Finland, known for hosting large areas of primary forests. Data completeness was lower for some central European countries. In the




Fig. 5 Workflow of the assessment of recent human disturbance in primary forest polygons.

case of Czechia, Slovakia, Poland and Romania, our data only accounted for 20–100% of the country-level estimates from Forest Europe¹⁰. For Austria, Switzerland and Hungary, instead, additional data on primary forests exists but it is not currently open-access, and therefore not considered here. The largest data gaps were in Sweden, Italy, Bulgaria, Estonia, Denmark and Russia, where our map accounted for less than 10% of the primary forest reported in Forest Europe¹⁰. The low data completeness found for Denmark likely depends on the inclusion of minimum-intervention forest reserves in Forest Europe (see⁴⁹) that were harvested until recently and therefore do not qualify as primary forests according to our definition.

Assessing recent human disturbance with remote sensing. Since our data were collected continuously over the last two decades, we cannot exclude that some forest patches may have undergone human disturbance after data collection. This is particularly relevant for areas where primary forests are lost at high rates, such as the Carpathians, Russian Karelia, or Northern Fennoscandia^{18–20}. To assess to what extent this might be an issue, we used the open-access Landsat archive and the LandTrendr disturbance detection algorithm^{50,51}, using Google Earth Engine⁵² (Fig. 5). Specifically, we 1) quantified the proportion of polygons in our map that underwent disturbance between 1985 and 2018, i.e., Landsat 5 operating time, 2) visually checked a stratified random selection of these disturbed polygons to quantify the prevalence of anthropogenic vs. natural disturbance, and 3) estimated the proportion of polygons in our map not meeting the necessary, but not sufficient, condition for being classified as primary (i.e. not being affected by anthropogenic disturbance within the last 35 years).

For each polygon contained in the map of primary forests, we extracted the whole stack of available Landsat images (~1985–today), and ran the LandTrendr⁵³ algorithm. LandTrendr identifies breakpoints in spectral time




Fig. 6 Examples of disturbed polygons, as detected by LandTrendr, before (left) and after (right) disturbance. **(a)** Natural disturbance in Babia Gora, Slovakia; **(b)** natural disturbance in the southern Bourgas Province of Bulgaria; **(c)** clear-cuts in Tatra National Park in Slovakia; **(d)** clearcuts in the Russian Republic of Karelia. Red circles are centred on the disturbed pixel randomly selected for visual inspection, and have a radius of 50 m; pink squares have a side of 1 km and were exclusively used to provide context reference to the photointerpreter. Image credits: Google Earth.

series, separates periods of disturbance or stability, and records the years in which disturbances occurred. To avoid problems due to cloud cover, changes in illumination, and atmospheric condition, we used all available images from the growing season of each year (1 May through 15 September) to derive yearly composite images⁵⁴. As our spectral index, we used Tasseled Cap Wetness (TCW), as this index is particularly sensitive to forest structure⁵⁵, is robust to spatial and temporal variations in canopy moisture⁵⁶, and consistently outperforms other spectral indices, including Normalized Difference Vegetation Index⁵³, for detecting forest disturbance^{50,57–59}. As input parameters for the LandTrendr algorithm when detecting forest disturbances, we used a prevalue of -300 TCW units, a minimum disturbance magnitude of 500 TCW units, and a maximum duration of 4 years.

After running LandTrendr, we eliminated noise by applying a minimum disturbance threshold (2 ha). We then visually inspected a stratified random selection of primary forest polygons highlighted as ‘disturbed’ by LandTrendr using very-high-resolution images available in Google Earth. For each biogeographic region, we randomly selected 20% of disturbed polygons up to a maximum of 100 polygons per region. Depending on the size of the polygons, we inspected up to 5 randomly selected disturbed pixels within each disturbed polygon with a minimum distance between pixels of 1 km. Based on the spectral and physical characteristics of the disturbed patch (brightness, shape, size), and on ancillary information derived from the Google Earth imagery, we assigned disturbance agents as either anthropogenic (i.e., forest harvest, infrastructure development) or natural (e.g., windstorm, bark beetle outbreak, fire; Figs. 6, 7). We conservatively considered a polygon as anthropogenically disturbed if at least a third of the points we checked for that polygon were anthropogenically disturbed. To avoid introducing an observer bias, all polygons were checked by the same photo-interpreter (FMS).

Out of the 17,309 polygons checked with LandTrendr, 4,734 (27.3% of total) experienced major disturbances between 1985 and 2018. The proportion of disturbed area was greater than 10% in 2,904 polygons. We visually inspected a total of 712 pixels across 268 primary forest polygons, corresponding to 1.5% of the total number of polygons and 5.7% of the disturbed polygons. We attributed a total of 149 pixels, across 61 primary forest polygons, to anthropogenic disturbance, i.e., 22.7% (bootstrapped standard error = 2.5%) of the polygons we checked (Table 2, Fig. 7). We thus estimated the total number of primary forest polygons being anthropogenically disturbed by multiplying the total number of polygons with the proportion of disturbed polygons (27.3%) and the share of these disturbed polygons attributed to anthropogenic causes (22.7%). This suggests our map contains 1,077 anthropogenically disturbed polygons (95% CIs [847, 1323]), which corresponds to 6.2% (95% CIs [4.9%, 7.6%]) of the total number of polygons. Disturbed polygons were concentrated in the Russian Federation (especially in Archangelsk region, Karelia and Komi republics), Southern Finland, and the Carpathians (Fig. 7; Table 2). The Boreal and Alpine biogeographical regions had the highest number of disturbed polygons (both




Fig. 7 Geographical distribution of naturally vs. anthropogenically disturbed polygons, as resulting from a visual check of 712 pixels across 268 polygons.

Biogeographic region	Num. PF polygons (1)	Num. disturbed PF polygons (2)	Num. disturbed PF polygons checked (3)	Num. of (3) with evident anthropogenic disturbance(4)	Share of (3) anthropogenically disturbed (4/3) %
Alpine	11,734	1,096	102	23	22.55
Arctic	96	105 [†]	20	0	0
Atlantic	83	48	13	0	0
Black Sea	19	6	1	0	0
Boreal	4,074	3,334	110	30	27.27
Continental	1,100	105	21	6	28.57
Macaronesia	27	8	2	1	50
Mediterranean	132	27	5	1	20
Pannonian	39	4	1	0	0
Steppic	5	1	0	0	0

Table 2. Recent human disturbance in primary forest polygons, summarized by biogeographical region. [†]The number of disturbed polygons is higher than the total number of polygons because some polygons expanding over more than one biogeographical region were split. PF – Primary Forest.

in total, and when considering only those with evident anthropogenic disturbance). The regions with the highest share of anthropogenically disturbed polygons were the Continental and Boreal region. The sample size in Macaronesia was too low to provide a reliable estimation of the incidence of human disturbance.

These estimates should be considered as lower bounds, because only the disturbance events with a magnitude sufficient to be captured with LandTrendr and occurring in 1985–2018 could be identified. Not being this a formal validation, the results presented here should not be extrapolated to primary forests not included in our map. Finally, being our database built with a bottom-up approach, we are unable to exclude the existence of remaining bias or interpretation error, which might have propagated through the successive steps required to build it. As such, we warn the users against possible heterogeneity in data quality, accuracy and completeness across datasets.

Usage Notes

All data files are referenced in a geographic coordinate system (lat/long, WGS 84 - EPSG code: 4326). The provided files are in a personal geodatabase, and can be accessed and displayed using standard GIS software such as: QGIS (www.qgis.org/en).

All datasets listed in Online-only Table 1 are freely available in Figshare (<https://doi.org/10.6084/m9.figshare.13194095.v1>)³³ with a Creative Commons CC BY 4.0 license. Two additional non open-access datasets are available on request to the corresponding author after approval of the respective copyright holders. These datasets are: ‘Hungarian Forest Reserve monitoring’ (ID 17, custodian: Ferenc Horváth); and ‘Potential OGF and primary forest in Austria’ (ID 48, custodian: Matthias Schickhofer). The same conditions apply for additional data

from the dataset ‘Strict Forest Reserves in Switzerland’ (ID 30, custodian: Jonas Stillhard). In the case of the dataset ‘Ancient and Primeval Beech Forests of the Carpathians and Other Regions of Europe’^{40,41} (ID 34, Custodian: UNESCO), this data is freely available online, but its copyright does not allow redistribution. We refer the interested reader to the website <https://www.protectedplanet.net/903141> for the original data.

Comments and requests of updates for the dataset are collected and discussed in the GitHub forum: <https://github.com/fmsabatini/PrimaryForestEurope>.

Code availability

The code to reproduce the composite layers, for post-processing and for assessing recent human disturbance with remote sensing is available together with the database in Figshare (<https://doi.org/10.6084/m9.figshare.13194095.v1>)³³. We included seven scripts:

- *00_ComposeMap.R* – Identifies overlapping polygons across individual datasets.
- *01_CreateComposite_Points.py* – Creates the composite point feature class.
- *02_CreateComposite_Polygons.py* – Creates the composite polygon feature class.
- *03_PostProcessing.R* – Extracts additional information on each primary forest.
- *04_Add_Postprocessing.py* – Imports post-processing output into the geodatabase.
- *05_Summary_stats.R* – Calculates summary statistics of primary forests
- *06_DisturbanceAssessment_Step1_exportIntermediateChangeImg.txt* – Runs LandTrendr in Google Earth Engine, tiles the area of interest, creates Change-Images for each tile, and exports these as intermediate .tif files containing the LandTrendr metrics.
- *07_DisturbanceAssessment_Step2_extractPolygonValuesFromChangeImg.txt* – Extracts LandTrendr metrics for each forest polygon from Change-Images and exports as .csv.

Python (.py) scripts were run in ESRI ArcGIS (v10.5) and are available also as ArcGIS Models inside the Geodatabase. R (.R) scripts were run using R (v 3.6.1)⁴⁶. The remaining .txt scripts were run in Google Earth Engine.

Received: 17 November 2020; Accepted: 6 July 2021;

Published: 17 August 2021

References

1. Watson, J. E. M. *et al.* The exceptional value of intact forest ecosystems. *Nat. Eco. Evo.* **2**, 599–610 (2018).
2. European Commission. in *COM(2020) 380 final* (Brussels, 2020).
3. Vandekerckhove, K. *et al.* Reappearance of Old-Growth Elements in Lowland Woodlands in Northern Belgium: Do the Associated Species Follow? *Silva Fenn.* **45**, 909–935 (2011).
4. Di Marco, M., Ferrier, S., Harwood, T. D., Hoskins, A. J. & Watson, J. E. Wilderness areas halve the extinction risk of terrestrial biodiversity. *Nature* **573**, 582–585 (2019).
5. Frey, S. J. K. *et al.* Spatial models reveal the microclimatic buffering capacity of old-growth forests. *Sci. Adv.* **2**, e1501392 (2016).
6. Zhou, G. Y. *et al.* Old-growth forests can accumulate carbon in soils. *Science* **314**, 1417–1417 (2006).
7. Burrascano, S., Keeton, W. S., Sabatini, F. M. & Blasi, C. Commonality and variability in the structural attributes of moist temperate old-growth forests: A global review. *For. Ecol. Manag.* **291**, 458–479 (2013).
8. Bauhus, J., Puettmann, K. & Messier, C. Silviculture for old-growth attributes. *For. Ecol. Manag.* **258**, 525–537 (2009).
9. Moore, K. D. In the shadow of the cedars: the spiritual values of old-growth forests. *Conserv. Biol.* **21**, 1120–1123 (2007).
10. FOREST EUROPE. State of Europe’s Forests 2015. (Ministerial Conference on the Protection of Forests in Europe, Madrid, 2015).
11. Ceccherini, G. *et al.* Abrupt increase in harvested forest area over Europe after 2015. *Nature* **583**, 72–77 (2020).
12. Levers, C. *et al.* Drivers of forest harvesting intensity patterns in Europe. *For. Ecol. Manag.* **315**, 160–172 (2014).
13. Potapov, P. *et al.* The last frontiers of wilderness: Tracking loss of intact forest landscapes from 2000 to 2013. *Sci. Adv.* **3** (2017).
14. Schickhofer, M. & Schwarz, U. *Inventory of Potential Primary and Old-Growth Forest Areas in Romania (PRIMOFARO). Identifying the largest intact forests in the temperate zone of the European Union.* (Euronatur Foundation, 2019).
15. Knorn, J. *et al.* Continued loss of temperate old-growth forests in the Romanian Carpathians despite an increasing protected area network. *Environ. Conserv.* **40**, 182–193 (2013).
16. Court of Justice of the European Union. C-441/17 - Commission v Poland (Forêt de Białowieża) *Judgment of the Court (Grand Chamber) of 17 April 2018* (2018).
17. Chylarecki, P. & Selva, N. Ancient forest: spare it from clearance. *Nature* **530**, 419–419 (2016).
18. Earthsight. Complicit in corruption. How billion-dollar firms and EU governments are failing Ukraine’s forests. (Earthsight, 2018).
19. Mikoláš, M. *et al.* Primary forest distribution and representation in a Central European landscape: Results of a large-scale field-based census. *For. Ecol. Manag.* **449**, 117466 (2019).
20. Hance, J. IKEA Logging Old-growth Forest for Low-price Furniture in Russia. <https://news.mongabay.com/2012/05/ikea-logging-old-growth-forest-for-low-price-furniture-in-russia/> (2012).
21. Sabatini, F. M. *et al.* Protection gaps and restoration opportunities for primary forests in Europe. *Divers. Distrib.* **26**, 1646–1662 (2020).
22. Barredo Cano, J. I. *et al.* *Mapping and assessment of primary and old-growth forests in Europe.* (EUR 30661 EN, Publications Office of the European Union, 2021).
23. Adam, D. & Vrška, T. *Important localities of old-growth forests in Landscape Atlas of the Czech Republic* (eds T Hrnčiarová, P Mackovčín, & I Zvara) (Ministry of Environment and Silva Tarouca Research Institute, Prague–Silva Tarouca Research Institute for Landscape and Ornamental Gardening, 2009).
24. Diaci, J. Virgin forests and forest reserves in Central and East European countries-History, present status and future development. *Proceedings of the invited lecturers’ reports presented at the COST E4 management committee and working groups meeting in Ljubljana, Slovenia* (1999).
25. Kirchmeir, H. & Kovarovic, A. *Nomination Dossier “Primeval Beech Forests of the Carpathians and Other Regions of Europe“ as extension to the existing Natural World Heritage Site “Primeval Beech Forests of the Carpathians and the Ancient Beech Forests of Germany” (1133bis).* (2016).
26. García Fedec, C., Berglund, H. & Strnad, M. Scoping document: information related to European old growth forests. (ETC/BD report to the EEA, 2015).

27. Veen, P. *et al.* Virgin forests in Romania and Bulgaria: results of two national inventory projects and their implications for protection. *Biodivers. Conserv.* **19**, 1805–1819 (2010).
28. Ibisch, P. L. & Ursu, A. Potential primary forests of Romania. (Greenpeace CEE Romania; Centre for Economics and Ecosystem Management, Eberswalde University for Sustainable Development; Geography Department, A. I. Cuza University of Iași, 2017).
29. Spracklen, B. D. & Spracklen, D. V. Identifying European Old-Growth Forests using Remote Sensing: A Study in the Ukrainian Carpathians. *Forests* **10**, 127 (2019).
30. Frank, G. *et al.* COST Action E27. *Protected Forest Areas in Europe-analysis and harmonisation (PROFOR): results, conclusions and recommendations.* (Federal Research and Training Centre for Forests, Natural Hazards and Landscape (BFW), 2007).
31. Sabatini, F. M. *et al.* Where are Europe's last primary forests? *Divers. Distrib.* **24**, 1426–1439 (2018).
32. McRoberts, R. E., Susanne, W., Gherardo, C. & Elizabeth, L. Assessing Forest Naturalness. *For. Sci.* **58**, 294–309 (2012).
33. Sabatini, F. M. *et al.* European Primary Forest Database. *figshare* <https://doi.org/10.6084/m9.figshare.13194095.v1> (2020).
34. FAO. *Global Forest Resources Assessment 2015. Terms and definitions.* (FAO, 2015).
35. Buchwald, E. *A hierarchical terminology for more or less natural forests in relation to sustainable management and biodiversity conservation in Proceedings: Third expert meeting on harmonizing forest-related definitions for use by various stakeholders* (Food and Agriculture Organization of the United Nations, 2005).
36. Blasi, C., Burrascano, S., Maturani, A. & Sabatini, F. M. *Old-growth forests in Italy.* (Palombi Editori, 2010).
37. Cateau, E. *et al.* *Le patrimoine forestier des réserves naturelles. Focus sur les forêts à caractère naturel. Cahier n°7.* (Réserves Naturelles de France, 2017).
38. Svoboda, M. *et al.* Landscape-level variability in historical disturbance in primary *Picea abies* mountain forests of the Eastern Carpathians, Romania. *J. Veg. Sci.* **25**, 386–401 (2014).
39. Potapov, P. *et al.* Mapping the world's intact forest landscapes by remote sensing. *Ecol. Soc.* **13** (2008).
40. Britz, H. *et al.* Nomination of the “Ancient Beech Forests of Germany” as Extension to the World Natural heritage “Primeval Beech Forests of the Carpathians”. *Nationale Naturlandschaften, Federal Republic of Germany. Nieden-stein: Specialised editing Cognitio Kommunikation & Planung* (2009).
41. UNEP-WCMC & IUCN. *Protected Area Profile for Ancient and Primeval Beech Forests of the Carpathians and Other Regions of Europe from the World Database of Protected Areas* <https://www.protectedplanet.net/903141> (2019).
42. EEA. *Biogeographical regions of Europe* <https://www.eea.europa.eu/data-and-maps/data/biogeographical-regions-europe-3> (2016).
43. EEA. European forest types. Categories and types for sustainable forest management reporting and policy. (EEA Technical Report No 9/2006. EEA, Copenhagen, 2006).
44. Bohn, U. *et al.* Map of the natural vegetation of Europe. Explanatory text with CD-ROM, (German Federal Agency for Nature Conservation, Bonn, Germany, 2003).
45. Hansen, M. C. *et al.* High-resolution global maps of 21st-century forest cover change. *Science* **342**, 850–853 (2013).
46. R Development Core Team. R: A language and environment for statistical computing v. 3.6.1. *R Foundation for Statistical Computing* <http://www.R-project.org/> (2019).
47. Miljødirektoratet. (2016).
48. FAO. Global Forest Resources Assessment 2015. Desk reference. 245 (FAO, Rome, 2015).
49. FOREST EUROPE. *Quantitative Indicators Country reports 2015* <https://foresteurope.org/state-europes-forests-2015-report/#1476295965372-d3bb1dd0-e9a0> (2015).
50. Cohen, W. B., Yang, Z., Healey, S. P., Kennedy, R. E. & Gorelick, N. A LandTrendr multispectral ensemble for forest disturbance detection. *Remote Sens. Environ.* **205**, 131–140 (2018).
51. Kennedy, E. R. *et al.* Implementation of the LandTrendr Algorithm on Google Earth Engine. *Remote Sensing* **10** (2018).
52. Gorelick, N. *et al.* Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sens. Environ.* **202**, 18–27 (2017).
53. Kennedy, R. E., Yang, Z. & Cohen, W. B. Detecting trends in forest disturbance and recovery using yearly Landsat time series: 1. LandTrendr—Temporal segmentation algorithms. *Remote Sens. Environ.* **114**, 2897–2910 (2010).
54. Griffiths, P., Van Der Linden, S., Kuemmerle, T. & Hostert, P. A pixel-based landsat compositing algorithm for large area land cover mapping *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* **6**, 2088–2101 (2013).
55. Cohen, W. B. & Spies, T. A. Estimating structural attributes of Douglas-fir/western hemlock forest stands from Landsat and SPOT imagery. *Remote Sens. Environ.* **41**, 1–17 (1992).
56. Czerwinski, C. J., King, D. J. & Mitchell, S. W. Mapping forest growth and decline in a temperate mixed forest using temporal trend analysis of Landsat imagery, 1987–2010. *Remote Sens. Environ.* **141**, 188–200 (2014).
57. Cohen, W. B., Yang, Z. & Kennedy, R. Detecting trends in forest disturbance and recovery using yearly Landsat time series: 2. TimeSync—Tools for calibration and validation. *Remote Sens. Environ.* **114**, 2911–2924 (2010).
58. Grogan, K., Pflugmacher, D., Hostert, P., Kennedy, R. & Fensholt, R. Cross-border forest disturbance and the role of natural rubber in mainland Southeast Asia using annual Landsat time series. *Remote Sens. Environ.* **169**, 438–453 (2015).
59. De Marzo, T. *et al.* Characterizing forest disturbances across the Argentine Dry Chaco based on Landsat time series. *International Journal of Applied Earth Observation and Geoinformation* **98**, 102310 (2021).
60. Frank, A. *Inventering av nyckelbiotoper: resultat till och med 2003.* (Skogsstyr., 2004).
61. Länsstyrelsen Västerbotten. *LstAC Skogar med höga naturvärden ovan gränsen för fjällnära skog 2003–2015* <https://ext-geodatakatalog.lansstyrelsen.se/GeodataKatalogen/> (2019).
62. Naturvårdsverket. *Skyddsvärda statliga skogar* <http://mdp.vic-metria.nu/miljodataportalen/GetMetaDataById?UID=3919E66E-2E09-440D-9171-B5074DF0C0ED> (2017).
63. Naturvårdsverket. *Skogliga värdekärnor* http://gpt.vic-metria.nu/data/land/skogliga_vardekarnor_2016.zip (2016).
64. Naturvårdsverket. *Preciserad kartering av kontinuitetsskog i Jämtlands län* http://gpt.vic-metria.nu/data/land/Preciserad_kskog_jamtland.zip (2019).
65. Ahlkrona, E., Giljam, C. & Wennberg, S. Kartering av kontinuitetsskogi boreal region. Metria AB på uppdrag av Naturvårdsverket (2017).
66. Naturvårdsverket. *Skyddad fjällbarrskog* https://gpt.vic-metria.nu/data/land/NMD/Skyddad_Fjallbarrskog.zip (2019).
67. Trotsiuk, V. *et al.* A mixed severity disturbance regime in the primary *Picea abies* (L.) Karst. forests of the Ukrainian Carpathians. *For. Ecol. Manag.* **334**, 144–153 (2014).
68. Kozák, D. *et al.* Profile of tree-related microhabitats in European primary beech-dominated forests. *For. Ecol. Manag.* **429**, 363–374 (2018).
69. Garbarino, M. *et al.* Gap disturbances and regeneration patterns in a Bosnian old-growth forest: a multispectral remote sensing and ground-based approach. *Ann. For. Sci.* **69**, 617–625 (2012).
70. Keren, S. *et al.* Comparative Structural Dynamics of the Janj Mixed Old-Growth Mountain Forest in Bosnia and Herzegovina: Are Conifers in a Long-Term Decline? *Forests* **5**, 1243–1266 (2014).
71. Motta, R. *et al.* Structure, spatio-temporal dynamics and disturbance regime of the mixed beech–silver fir–Norway spruce old-growth forest of Biogradska Gora (Montenegro). *Plant Biosyst.* **149**, 966–975 (2015).
72. Motta, R. *et al.* Development of old-growth characteristics in uneven-aged forests of the Italian Alps. *Eur. J. For. Res.* **134**, 19–31 (2015).
73. Panayotov, M. *et al.* *Mountain coniferous forests in Bulgaria – structure and natural dynamics.* (University of Forestry and Geosoft, 2016).

74. Lõhmus, A. & Kraut, A. Stand structure of hemiboreal old-growth forests: Characteristic features, variation among site types, and a comparison with FSC-certified mature stands in Estonia. *For. Ecol. Manag.* **260**, 155–165 (2010).
75. EEA. *Developing a forest naturalness indicator for Europe. Concept and methodology for a high nature value (HNV) forest indicator.* (EEA Technical report No 13/2014, Luxembourg: Publications Office of the European Union, 2014).
76. Rossi, M., Bardin, P., Cateau, E. & Vallauri, D. Forêts anciennes de Méditerranée et des montagnes limitrophes: références pour la naturalité régionale. *WWF France, Marseille, France*, 144 (2013).
77. Myhre, T. Skogkur 2020. redningsplan for Norges unike skoger. *WWF Verdens villmarksfond, Norges naturvernforbund, SABIMA* (2012).
78. Ruete, A., Snäll, T. & Jönsson, M. Dynamic anthropogenic edge effects on the distribution and diversity of fungi in fragmented old-growth forests. *Ecol. Appl.* **26**, 1475–1485 (2016).
79. Heiri, C., Wolf, A., Rohrer, L., Brang, P. & Bugmann, H. Successional pathways in Swiss mountain forest reserves. *Eur. J. For. Res.* **131**, 503–518 (2012).
80. Brang, P., Heiri, C. & Bugmann, H. *Waldreservate: 50 Jahre natürliche Waldentwicklung in der Schweiz.* (Haupt, 2011).
81. Pantić, D. *et al.* Structural, production and dynamic characteristics of the strict forest reserve Račanska šljivovica on Mt. Tara. *Glasnik Šumarskog fakulteta*, 93–114 (2011).
82. Savoie, J. M. *et al.* Vieilles forêts pyrénéennes de Midi-Pyrénées. Deuxième phase. *Evaluation et cartographie des sites. Recommandations. Rapport final.* (Ecole d'Ingénieurs de PURPAN/DREAL Midi-Pyrénées, 2015).
83. Savoie, J. M. *et al.* Forêts pyrénéennes anciennes de Midi-Pyrénées. Rapport d'Etude de projet FEDER 2008–2011. 320 (Ecole d'Ingénieurs de PURPAN/DREAL Midi-Pyrénées, 2011).
84. WWF Finland. Kansallisomaisuus turvaan - valtion omistamia suojelemaan metsä- ja suoalueita, (WWF Suomen raportteja, 2012).
85. Kitenberga, M. *et al.* A mixture of human and climatic effects shapes the 250-year long fire history of a semi-natural pine dominated landscape of Northern Latvia. *For. Ecol. Manag.* **441**, 192–201 (2019).
86. Baders, E., Senhofa, S., Purina, L. & Jansons, A. Natural succession of Norway spruce stands in hemiboreal forests: case study in Slitere national park, Latvia. *Baltic Forestry* **23**, 522–528 (2017).
87. Kokarēviča, I. *et al.* Vegetation changes in boreo-nemoral forest stands depending on soil factors and past land use during an 80 year period of no human impact. *Can. J. For. Res.* **46**, 376–386 (2016).
88. Fernandez López, A. B. Parque Nacional de Garajonay, Patrimonio Mundial. (Organismo Autonomo Parques Nacionales, 2009).
89. TRAGSATEC. Segundo inventario ecológico del Parque Nacional de Garajonay. (Parque Nacional de Garajonay, 2006).
90. Fernández, A. B. & Gómez, L. Qué son los bosques antiguos de laurisilva. Su valor y situación en Canarias. *La Gomera, entre bosques y taparuchas*, 177–236 (2016).
91. Matović, B. *et al.* Comparison of stand structure in managed and virgin European beech forests in Serbia. *Šumarski list* **142**, 47–57 (2018).
92. Kiš, A., Stojišić, V., & Dinić, A. In *2nd International Symposium on Nature Conservation. Proceedings* 373–382 (Institute for Nature Conservation of Vojvodina Province, Novi Sad, 2016).
93. Kobayakov, K. & Jakolev, J. Atlas of high conservation value areas, and analysis of gaps and representativeness of the protected area network in northwest Russia. (Finnish Environment Institute, 2013).
94. Diku, A. & Shuka, L. *Pyjet e vjetër të ahut në shqipëri (Old Beech forests in Albania).* (PSEDA - ILIRIA, 2017).
95. Burrascano, S. *et al.* It's a long way to the top: Plant species diversity in the transition from managed to old-growth forests. *J. Veg. Sci.* **29**, 98–109 (2018).

Acknowledgements

This study was realized and funded by the project “Policy and on-ground action for primary forest protection, boreal and temperate primary forests” funded through the Griffith University (Australia) and implemented by the Frankfurt Zoological Society (FZS) and Wild Europe Initiative as well as the Naturwald Akademie. Additional funding derive from the European Commission (Marie Skłodowska-Curie fellowship to FMS, project FORESTS & CO, #658876). The Italian dataset was supported by funding from the Department for Nature Protection of the Italian Ministry of the Environment, Land and Sea Protection. The GEVFP's French dataset was supported by funding from EU, RF and ‘Conseil Régional Midi-Pyrénées’. MM gratefully acknowledges the Czech Science Foundation (Grant GACR no. 21-27454S), as well as the institutional project EVA No.CZ.02.1.01/0.0/0.0/16_019/0000803. The Serbian dataset was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia, grant number 451-03-9/2021-14/ 200197. This work would not have been possible without all those who responded to the questionnaires and those who collected all the data presented here. A special thanks goes to all data contributors, especially those who are not coauthors of this paper (in alphabetic order): Paloma Hannonen (Dataset 39), Matti Liimatainen (Dataset 38), Simone Mayrhofer (Dataset 49), Daniel Vallauri (Dataset 13) and Juraj Vysoky (Dataset 26). FMS also acknowledges the financial support within the funding programme Open Access Publishing by the German Research Foundation (DFG). We also thank Teresa De Marzo for support during the assessment of recent human disturbance in primary forest polygons. We thank the editor and two anonymous reviewers for very helpful and constructive comments on prior manuscript versions.

Author contributions

The original idea for the database is from F.M.S., T.K. and Z.K., F.M.S. and H.B. harmonized the datasets, and conducted the literature review. D.A., J.A.A., E.B., S.B., E.C., A.D., I.M.D., A.B.F., M.G., N.G., F.H., S.K., M.K., A.K.i., A.K.r., P.L.I., L.L., F.L., B.M., R.N.M., P.M., S.M., R.M., M.M., G.M., M.P., R.P., L.N., A.R., M.S., B.S., J.S., D.S., J.S., O.-P.T., E.T., R.V., T.V., M.W., M.Y., T.Z., A.Z. contributed data. F.M.S., H.B. and T.K. created the first draft of the manuscript and all co-authors contributed substantially to its revision.

Funding

Open Access funding enabled and organized by Projekt DEAL.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to F.M.S.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

The Creative Commons Public Domain Dedication waiver <http://creativecommons.org/publicdomain/zero/1.0/> applies to the metadata files associated with this article.

© The Author(s) 2021, corrected publication 2021

Francesco Maria Sabatini^{1,2}✉, Hendrik Bluhm³, Zoltan Kun⁴, Dmitry Aksenov⁵, José A. Atauri⁶, Erik Buchwald⁷, Sabina Burrascano⁸, Eugénie Cateau⁹, Abdulla Diku¹⁰, Inês Marques Duarte¹¹, Ángel B. Fernández López¹², Matteo Garbarino¹³, Nikolaos Grigoriadis¹⁴, Ferenc Horváth¹⁵, Srđan Keren¹⁶, Mara Kitenberga¹⁷, Alen Kiš¹⁸, Ann Kraut¹⁹, Pierre L. Ibisich²⁰, Laurent Larrieu^{21,22}, Fabio Lombardi²³, Bratislav Matovic²⁴, Radu Nicolae Melu²⁵, Peter Meyer²⁶, Rein Midteng²⁷, Stjepan Mikac²⁸, Martin Mikoláš^{29,30}, Gintautas Mozgeris³¹, Momchil Panayotov³², Rok Pisek³³, Leónia Nunes³⁴, Alejandro Ruete³⁵, Matthias Schickhofer³⁶, Bojan Simovski³⁷, Jonas Stillhard³⁸, Dejan Stojanovic³⁹, Jerzy Szwagrzyk⁴⁰, Olli-Pekka Tikkanen⁴¹, Elvin Toromani⁴², Roman Volosyanchuk^{43,44}, Tomáš Vrška⁴⁵, Marcus Waldherr⁴⁶, Maxim Yermokhin⁴⁷, Tzvetan Zlatanov⁴⁸, Asiya Zagidullina⁴⁹ & Tobias Kuemmerle⁵⁰

¹German Centre for Integrative Biodiversity Research (iDiv) - Halle-Jena-Leipzig, Puschstrasse 4, 04103, Leipzig, Germany. ²Martin-Luther-Universität Halle-Wittenberg, Institut für Biologie, Am Kirchtor 1, 06108, Halle, Germany. ³Humboldt-Universität zu Berlin, Geography Department, Unter den Linden 6, 10099, Berlin, Germany. ⁴Frankfurt Zoological Society, Bernhard-Grzimek-Allee 1, 60316, Frankfurt, Germany. ⁵NGO "Transparent World", Rossolimo str. 5/22, building 1, 119021, Moscow, Russia. ⁶EUROPARC-Spain/Fundación Fernando González Bernádez. ICEI Edificio A. Campus de Somosaguas, E28224, Pozuelo de Alarcón, Spain. ⁷The Danish Nature Agency, Gjøddinggård, Førstballevej 2, DK-7183, Randbøl, Denmark. ⁸Sapienza University of Rome, Department of Environmental Biology, P.le Aldo Moro 5, 00185, Rome, Italy. ⁹Réserves Naturelles de France, La Bourdonnerie, Dijon cedex, 21000, France. ¹⁰PSEDA-ILIRIA. Forestry department, Tirana, 1000, Albania. ¹¹Centre for Applied Ecology "Professor Baeta Neves" (CEABN), InBIO, School of Agriculture, University of Lisbon, Tapada da Ajuda, 1349-017, Lisbon, Portugal. ¹²Parque Nacional de Garajonay. Avda. V Centenario, edif. Las Creces, local 1, portal 3, 38800 San Sebastian de La Gomera, Tenerife, Spain. ¹³University of Torino, Department DISAFA L.go Paolo Braccini 2, Grugliasco, 10095, Italy. ¹⁴Forest Research Institute, Vassilika, 57006, Thessaloniki, Greece. ¹⁵Centre for Ecological Research, Institute of Ecology and Botany, Alkotmány u. 2-4., 2163, Vácrátót, Hungary. ¹⁶Faculty of Forestry, University of Agriculture in Krakow, aleja 29-Listopada 46, 31-415, Krakow, Poland. ¹⁷Latvian State Forest Research Institute "Silava", Rigas street 111, Salaspils, LV-2169, Latvia. ¹⁸Institute for Nature Conservation of Vojvodina Province, Radnička 20a, Novi Sad, 21000, Serbia. ¹⁹University of Tartu, Institute of Ecology and Earth Sciences, Vanemuise 46, EE-51014, Tartu, Estonia. ²⁰Centre for Ecomics and Ecosystem Management, Faculty of Forest and Environment, Eberswalde University for Sustainable Development, Alfred-Möller-Str. 1, 16225, Eberswalde, Germany. ²¹Université de Toulouse, INRAE, UMR DYNFOR, 24 Chemin de Borde-Rouge Auzeville CS 52627, Castanet-Tolosan, 31326, France. ²²CRPF-Occitanie, antenne de Tarbes, place du foirail, 65000, Tarbes, France. ²³Mediterranean University of Reggio Calabria, Agraria Department, Loc. Feo di Vito, 89122, Reggio Calabria, Italy. ²⁴University of Novi Sad, Institute of Lowland Forestry and Environment, Antona Cehova 13d, Novi Sad, 21102, Serbia. ²⁵World Wide Fund for nature (CEE), Lunga street 190, Brasov, 500051, Romania. ²⁶Northwest German Forest Research Institute, Department Forest Nature Conservation, Professor-Oelkers-Straße 6, 34346. Hann, Münden, Germany. ²⁷Asplan Viak A.S.Kjörboveien 20, postboks 24, N-1300, Sandvika, Norway. ²⁸University of Zagreb, Faculty of Forestry, Svetosimunska cesta 25, 10000, Zagreb, Croatia. ²⁹Czech University of Life Sciences, Faculty of Forestry and Wood Sciences, Kamýcka cesta 1176, CZ-16521, Praha6-Suchbát, Czech Republic. ³⁰PRALES, Odrnovie 563, SK-01322, Rosina, Slovakia. ³¹Vytautas Magnus University, K. Donelaičio g. 58, LT-44248, Kaunas, Lithuania. ³²University of Forestry, Dendrology Department, bulvard "Sveti Kliment Ohridski" 10, 1756, Sofia, Bulgaria. ³³Slovenia Forest

Service, Department for forest management planning, Vecna pot 2, 1000, Ljubljana, Slovenia. ³⁴Centre for Applied Ecology "Professor Baeta Neves" (CEABN), InBIO, School of Agriculture, University of Lisbon, Tapada da Ajuda 1349-017, Lisbon, Portugal. ³⁵Greensway AB, Ulls väg 24A. 756 51, Uppsala, Sweden. ³⁶Freelance forest expert and book author, Vienna, Austria. ³⁷Ss. Cyril and Methodius University in Skopje, Hans Em Faculty of Forest Sciences, Landscape Architecture and Environmental Engineering, Department of Botany and Dendrology, P.O. Box 235, MK-1000, Skopje, North Macedonia. ³⁸Swiss Federal Research Institute for Forest, Snow and Landscape Research WSL, Forest Resources and Management, Zürcherstrasse 111, 8903, Birmensdorf, Switzerland. ³⁹University of Novi Sad, Institute of Lowland Forestry and Environment, Antona Cehova 13d, Novi Sad, 21000, Serbia. ⁴⁰Department of Forest Biodiversity, University of Agriculture, Kraków, Poland. ⁴¹University of Eastern Finland, School of forest Sciences, Yliopistokatu 7, 80100, Joensuu, Finland. ⁴²Agricultural University of Tirana, Forestry Department, Kodër Kamëz, SH1, 1029, Tirana, Albania. ⁴³World Wide Fund for nature (DCP) Ukraine, Mushaka 48, Lviv, 79011, Ukraine. ⁴⁴Ecosphera NGO, Kapushans'ka 82a, Uzhhorod, 88000, Ukraine. ⁴⁵Silva Tarouca Research Institute, Department of Forest Ecology, Lidická 25/27, 602 00, Brno, Czech Republic. ⁴⁶Centre for Ecnics and Ecosystem Management, Faculty of Forest and Environment, Eberswalde University for Sustainable Development, Alfred-Möller-Str. 1, 16225, Eberswalde, Germany. ⁴⁷Institute of Experimental Botany of the National Academy of Sciences of Belarus, Laboratory of Productivity & Stability of Plant Communities, 220072, Akademicheskaya St. 27, Minsk, Belarus. ⁴⁸Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, 2 Gagarin Street, 1113, Sofia, Bulgaria. ⁴⁹Saint-Petersburg State University, Department of Vegetation Science, University Embankment, 7/9, St Petersburg, 199034, Russia. ⁵⁰Humboldt-Universität zu Berlin, Geography Department & Integrative Research Institute on Transformation in Human-Environment Systems, Unter den Linden 6, 10099, Berlin, Germany. ✉e-mail: francesco.sabatini@botanik.uni-halle.de